

04-03-00

A

03/31/00
 1c644 U.S. PTO

UTILITY PATENT APPLICATION TRANSMITTAL <i>(Only for new nonprovisional applications under 37 CFR 1.53(b))</i>	Attorney Docket No.	B0801/7170 (JRV)
	First Named Inventor or Application Identifier	
	Kupper et al.	
	Express Mail Label No.	EL018098005US
Date of Deposit		March 31, 2000

APPLICATION ELEMENTS <i>See MPEP chapter 600 concerning utility patent application contents</i>	ADDRESS TO: Box Patent Application Assistant Commissioner for Patents Washington, DC 20231
1. <input type="checkbox"/> Fee Transmittal Form <i>(Submit an original, and a duplicate for fee processing)</i> 2. <input checked="" type="checkbox"/> Specification [Total pages 36] 29 - pages specification 1 - pages abstract 6 - pages claims 50 - Total claims 3. <input checked="" type="checkbox"/> Drawing(s) (35 USC 113) [Total sheets 1] <input type="checkbox"/> Informal <input checked="" type="checkbox"/> Formal [Total drawings 1] 4. <input type="checkbox"/> Oath or Declaration [Total pages] a. <input type="checkbox"/> Newly executed (original or copy) b. <input type="checkbox"/> Unsigned c. <input type="checkbox"/> Copy from a prior application (37 CFR 1.63(d)) <i>(for continuation/divisional with Box 17 completed)</i> [Note Box 5 below] i. <input type="checkbox"/> DELETION OF INVENTOR(S) Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) and 1.33(b). 5. <input type="checkbox"/> Incorporation by Reference <i>(usable if Box 4c is checked)</i> The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4c, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.	6. <input type="checkbox"/> Microfiche Computer Program (Appendix) 7. <input type="checkbox"/> Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary) a. <input type="checkbox"/> Computer Readable Copy b. <input type="checkbox"/> Paper Copy (identical to computer copy) c. <input type="checkbox"/> Statement verifying identity of above copies ACCOMPANYING APPLICATION PARTS 8. <input type="checkbox"/> Assignment Papers/cover sheet & documents(s) 9. <input type="checkbox"/> 37 CFR 3.73(b) Statement <i>(when there is an assignee)</i> <input type="checkbox"/> Power of Attorney 10. <input type="checkbox"/> English Translation of Document <i>(if applicable)</i> 11. <input type="checkbox"/> Information Disclosure Statement PTO-1449 <input type="checkbox"/> Copies of IDS Citations 12. <input type="checkbox"/> Preliminary Amendment 13. <input checked="" type="checkbox"/> Return Receipt Postcard (MPEP 503) <i>(Should be specifically itemized)</i> 14. <input type="checkbox"/> Small Entity Statement(s) <input type="checkbox"/> Statement filed in prior application, Status still proper and desired 15. <input type="checkbox"/> Certified Copy of Priority Document(s) <i>(if foreign priority is claimed)</i>
16. Other:	
17. If a CONTINUING APPLICATION , check appropriate box and supply the requisite information: <input type="checkbox"/> Continuation <input type="checkbox"/> Divisional <input type="checkbox"/> Continuation-in-part (CIP) of prior application No.: <input type="checkbox"/> Cancel in this application original claims of the prior application before calculating the filing fee. <input type="checkbox"/> Amend the specification by inserting before the first line the sentence: This application is a <input type="checkbox"/> continuation <input type="checkbox"/> divisional of application serial no.	

18. CORRESPONDENCE ADDRESS

Correspondence address below

ATTORNEY'S NAME	John R. Van Amsterdam, Reg. No. 40,212				
NAME	Wolf, Greenfield & Sacks, P.C.				
ADDRESS	600 Atlantic Avenue				
CITY	Boston	STATE	MA	ZIP	02210
COUNTRY	USA	TELEPHONE	(617) 720-3500	FAX	(617) 720-2441

19. SIGNATURE OF APPLICANT, ATTORNEY, OR AGENT REQUIRED

NAME	John R. Van Amsterdam, Reg. No. 40,212
SIGNATURE	<i>John R. Van Amsterdam</i>
DATE	<i>March 31, 2000</i>

- 1 -

MODIFIED DENDRITIC CELLS AND USES THEREFOR

Government Support

This work was funded in part by the National Institutes of Health under grant numbers
5 AI25082-11, AR42689 and HL54936. The government may retain certain rights in this
invention .

Related Applications

This application claims the benefit of United States provisional application serial no.
10 60/127,423, filed April 1, 1999.

Field of the Invention

The invention relates to antigen presenting cells which are modified with exogenous
binding molecules which bind selectin ligands on endothelial cells.

Background of the Invention

The immune system of a mammal often provides the first line of defense against
pathogenic organisms, as well as against tumors. The immune system recognizes antigens
expressed by tumor cells or pathogens as foreign, i.e., "non-self". Upon recognition of a non-
15 self antigen, an immune response is mounted against the antigen, resulting in antibodies
and/or cytolytic T cells which recognize the antigen. The immune response of a mammal is
20 also responsible for allergy (to antigens known as allergens) and autoimmune disease, which
results from inappropriate recognition of host proteins as non-self.

The process by which the mammalian immune system recognizes and reacts to foreign
25 or alien materials is a complex one. An important facet of the system is the T cell response.
This response requires that T cells recognize and interact with complexes of cell surface
molecules, referred to as human leukocyte antigens ("HLA"), or major histocompatibility
complexes ("MHCs"), and peptides. The peptides are derived from larger molecules which
are processed by the cells which also present the HLA/MHC molecule. The interaction of T
30 cells and complexes of HLA/peptide is restricted, requiring a T cell specific for a particular
combination of an HLA molecule and a peptide. Another mechanism exists for presentation
of non-peptide antigens (such as lipids, glycolipids and carbohydrates) to T cells. This

mechanism involves the formation of a complex of the non-peptide antigen and CD1 proteins.

If a specific T cell is not present, there is no T cell response even if its partner complex is present. Similarly, there is no response if the specific complex is absent, but the T cell is present. These mechanisms are involved in the immune system's response to foreign materials, in autoimmune pathologies, and in responses to cellular abnormalities.

It has been demonstrated that cytotoxic T lymphocytes provide an effective response against tumor cells. Tumor cells express tumor associated genes. The protein expression products of these genes are processed into peptides which, in turn, are presented by HLA/MHC on cell surfaces, which can lead to lysis of the tumor cells by specific CTLs. The genes are said to code for proteins which are "tumor rejection antigen precursors", and the peptides derived therefrom are referred to as "tumor rejection antigens". Therefore, immunization which increases host cytotoxic T lymphocytes specific for one or more tumor rejection antigens can reduce tumor load in the host. CTL involvement in other desirable immune responses (e.g. against antigens of pathogens) as well as in undesirable immune responses (e.g. against allergens, self antigens and antigens associated with transplanted tissue grafts) is also well known.

Several steps are required for the generation of a specific T cell response, starting with antigen processing and presentation of antigenic peptides by MHC on antigen presenting cells (APCs), followed by T cell priming from signals transmitted through the T cell receptor and through co-stimulatory molecules, such as CD28 and CD40 ligand, resulting in the clonal expansion of T cells.

Dendritic cells (DCs) are potent antigen presenting cells which efficiently stimulate immune recognition of presented peptides. Activation of naive T cells in lymph nodes is mediated exclusively by dendritic cells. A major weakness of current DC immunization approaches is the inability to deliver the DCs directly from blood to secondary lymphoid tissues (including lymph nodes) throughout the body where selectin ligands on endothelial cells are required for homing of blood-borne leukocytes. Thus there is a need to have improved methods and compositions for stimulating immune responses against pathogenic organisms, cellular abnormalities including tumors, and the like.

There is also a need to have improved methods and compositions for reducing inappropriate immune responses, such as found in allergy and autoimmune disease. Reduction of host immune responses is also desirable to reduce allograft transplant rejection

by the host.

Summary of the Invention

It has now been discovered that dendritic cells cultured *in vitro* do not express sufficient L-selectin, a critical adhesion molecule that permits entry of cells through the lymph node post capillary venule endothelium, for effective targeting of dendritic cells to peripheral lymph nodes. Cells exiting blood into lymph nodes pass through high endothelial venules (HEVs), which are specialized blood vessels which constitutively express carbohydrate-containing L-selectin ligands. For example, naive T cells express L-selectin on their surface which allows them to interact with L-selectin ligands on HEV to bind under flow conditions. The bound T cells then are activated by chemokines *in situ* and use cell surface integrins to extravasate into lymph nodes. Because they lack sufficient L-selectin expression, cultured dendritic cells cannot exit blood via HEVs directly into lymph node where naive T cells await antigen presentation.

To remedy this deficiency, isolated dendritic cells have been genetically modified to express on their cell surface molecules which bind to L-selectin ligands present on HEVs. Dendritic cells also have been treated with activated platelets which express P-selectin to form platelet modified dendritic cells. The activated platelets serve as a bridge between the cultured dendritic cells and selectin ligands on HEVs. Both genetically modified and platelet modified DCs adhere to lymph node vessel specific ligands *in vitro* and *in vivo*.

Thus the invention provides isolated genetically modified dendritic cells, optionally treated with activated platelets or membrane microparticles thereof. The invention also provides isolated platelet modified dendritic cells. Methods for delivering the modified dendritic cells to peripheral lymph nodes and methods for using the modified dendritic cells to stimulate immune responses also are provided. Vaccine compositions containing the modified dendritic cells also are provided.

According to one aspect of the invention, methods for delivery of dendritic cells to a secondary lymphoid tissue of a subject or to a non-lymphoid tissue of a subject where selectin ligands are expressed on endothelial cells are provided. The methods include providing isolated genetically modified dendritic cells which express on the cell surface a selectin polypeptide which includes an endothelial selectin ligand binding portion of a selectin. The binding portion of a selectin is a portion of L-selectin, E-selectin or P-selectin. The methods

also include administering the isolated genetically modified dendritic cells to the subject. Preferably the selectin polypeptide is L-selectin, E-selectin or P-selectin.

In certain preferred embodiments, the step of providing isolated dendritic cells includes isolating dendritic cells from the subject and transfecting the isolated dendritic cells with a nucleic acid molecule that encodes the selectin polypeptide. The nucleic acid molecule preferably is an expression vector. Preferred expression vectors included retroviruses, lentiviruses, adenoviruses and lambda bacteriophages. A particularly preferred expression vector is a retrovirus.

The step of providing isolated dendritic cells also can include treating the isolated transfected dendritic cells with isolated activated platelets or membrane microparticles thereof which contain P selectin.

According to another aspect of the invention, methods for delivery of dendritic cells to a secondary lymphoid tissue of a subject or to a non-lymphoid tissue of a subject where selectin ligands are expressed on endothelial cells are provided. The methods include providing isolated dendritic cells, treating the isolated dendritic cells with isolated activated platelets or membrane microparticles thereof which contain P selectin to form platelet modified dendritic cells, and administering the isolated platelet modified dendritic cells to the subject. In some embodiments, the step of providing isolated dendritic cells further comprises culturing the isolated dendritic cells to expand the isolated dendritic cells. In another aspect of the invention, the isolated dendritic cells and the isolated activated platelets or membrane microparticles thereof are not contacted, but the isolated activated platelets or membrane microparticles thereof are administered prior to or concurrently with the isolated dendritic cells.

In certain preferred embodiments of the foregoing methods, the secondary lymphoid tissue is peripheral lymph nodes, appendix or tonsil. In other embodiments, the non-lymphoid tissue where selectin ligands are expressed on endothelial cells is a site of chronic inflammation.

In other preferred embodiments of the foregoing methods, the step of providing isolated dendritic cells further comprises culturing the isolated dendritic cells to expand the isolated dendritic cells.

Preferably the isolated dendritic cells are administered parenterally, especially intraarterially or intravenously.

In the foregoing methods, the step of providing isolated dendritic cells in some embodiments includes culturing the isolated dendritic cells to expand the isolated dendritic cells.

According to still another aspect of the invention, compositions including isolated genetically modified dendritic cells are provided. The dendritic cells express on the cell surface a selectin polypeptide having an endothelial selectin ligand binding portion of L-selectin, E-selectin or P-selectin. Preferred selectin polypeptides are L-selectin, E-selectin and P-selectin.

In preferred embodiments, the amount of the selectin polypeptide expressed on the cell surface is greater than the naturally occurring amount of the selectin expressed on the cell surface *in vitro*. Preferably the amount of the selectin polypeptide expressed is sufficient to target the genetically modified dendritic cells to secondary lymphoid tissues, including peripheral lymph nodes, or non-lymphoid tissues having selectin ligands expressed on endothelial cells.

In some embodiments the isolated dendritic cells are transfected with a nucleic acid molecule which encodes the selectin polypeptide, and preferably the nucleic acid molecule is an expression vector. Preferred expression vectors include retroviruses, lentiviruses, adenoviruses and bacteriophage lambda, particularly retroviruses.

In other embodiments, the compositions include isolated activated platelets or membrane microparticles thereof which contain P selectin.

According to another aspect of the invention, compositions including isolated dendritic cells and isolated activated platelets or membrane microparticles thereof which contain P selectin are provided.

Vaccine compositions including the foregoing compositions and an antigen also are provided in another aspect of the invention. In certain preferred embodiments, the antigen is a peptide. In other preferred embodiments, the isolated dendritic cells are loaded with the antigen *ex vivo*, or the isolated dendritic cells are transfected with a nucleic acid molecule which encodes the antigen. In these latter embodiments, the nucleic acid molecule preferably encodes at least two antigens. Preferred antigens include MAGE, MART, LAGE, NY-ESO-1, tyrosinase, PRAME, prostate specific antigen (PSA), BCR-ABL, T cell receptor from T cell tumors, immunoglobulins from B cell tumors, RAS and carcinoembryonic antigen (CEA).

In still other preferred embodiments, the vaccine compositions include at least one

adjuvant. In some of these preferred embodiments, the adjuvant is a nucleic acid molecule encoding a polypeptide adjuvant transfected into the isolated dendritic cells. Certain preferred adjuvants stimulate an immune response to the antigen, including IL-12, GM-CSF, CD40, CD80 (B7-1) and CD86 (B7-2). Other preferred adjuvants inhibit an immune response to the antigen, including IL-10 and TGF- β . Still other preferred adjuvants orient an immune response to certain lymphocyte subsets, including IL-4 (Th2) and interferon- γ (Th1).

According to yet another aspect of the invention, methods for stimulating or inhibiting an immune response to an antigen in a subject are provided. The methods include administering to the subject an appropriate composition selected from the foregoing vaccine compositions.

Methods for testing the immunogenicity of an antigen also are provided. The methods include providing a subject, administering to the subject one of the foregoing vaccine composition (except those vaccine compositions which inhibit an immune response), wherein the immunogenicity of the antigen is not known, and determining the immune response of the subject to the antigen.

Also provided are methods for making the compositions and vaccine compositions described herein. The methods include in some embodiments isolating dendritic cells and introducing into the dendritic cells a selectin polypeptide, an antigen, a cytokine, an adjuvant, a nucleic acid molecule which encodes any of the foregoing, or combinations of the foregoing.

Kits which include the compositions also are provided, optionally including instructions for administering the compositions to modulate an immune response. The kits can also include other components, such as adjuvants, cytokines, antigens, and the like.

Still other aspects of the invention provide for the use of the foregoing compositions in the preparation of a medicament. Preferred medicaments include those which are useful in modulation of an immune response in a subject.

These and other objects of the invention will be described in further detail in connection with the detailed description of the invention.

Brief Description of the Drawings

Fig. 1 is a digitized low-voltage scanning electron micrograph showing delineating E/L-selectin expression on the surface of a murine bone marrow-derived dendritic cell at

80,000 x magnification.

Detailed Description of the Invention

It has now been discovered that cultured dendritic cells (DCs) can be modified to enable the DCs to enter peripheral lymph nodes by extravasation through high endothelial venules (HEVs). Although not wishing to be bound by any particular theory or mechanism, it is believed that a principal reason why cultured DCs are unable to reach peripheral lymph nodes is because cultured DCs lack sufficient L-selectin expression. This unexpected finding provides the basis for designing methods for overcoming this deficiency, namely, by genetically modifying isolated dendritic cells to express a selectin polypeptide which binds a selectin ligand on endothelial cells. Alternatively, entry of cultured DCs into peripheral lymph node (PLN) also can be made possible by treating isolated dendritic cells with activated platelets or a derivative thereof which includes P-selectin protruding from a surface where it is accessible for binding by dendritic cells and endothelial selectin ligand on HEVs. According to yet another aspect of the invention, DCs also can be both genetically modified and treated with activated platelets or derivatives thereof. Therefore the invention disclosed herein provides the unexpected result that augmentation of selectin polypeptides on the surface of cultured dendritic cells can alter the ability of the DCs to enter peripheral lymph nodes, thereby enhancing antigen presentation. Thus the invention provides methods for delivering or targeting antigen presenting cells to peripheral lymph nodes, methods for modulating an immune response to an antigen presented by dendritic cells, and compositions useful therefor.

As used herein, a "selectin polypeptide" is a polypeptide which includes a portion of a selectin molecule which is capable of binding selectively to a selectin ligand on endothelial cells. Such selectin polypeptides include L-selectin, E-selectin, P-selectin, including non-cleavable forms of L-, E- and P- selectins, such as those having mutated protease cleavage sites (see, e.g., Stoddart et al., *J. Immunol.* 157:5653-5659, 1996), chimeric selectin molecules, e.g. E/L-selectin (von Andrian et al., *Cell* 82:989-999, 1995; Stein et al. *J. Exp. Med.* 189:37-49, 1999), and functional variants of the foregoing which retain selectin binding properties.

As used herein, "selectin ligands on endothelial cells" include molecules present on the surface of high endothelial venules which bind L-selectin. Selectin ligands on endothelial

cells in some instances are known as peripheral node addressins (PNADs). L-selectin recognizes by selective binding several glycoprotein ligands, including GlyCAM-1 (Sgp50), CD34 (Sgp-90), and Sgp200, present on the high endothelial venules of the peripheral lymph nodes.

As used herein, an "endothelial selectin ligand binding portion of a selectin" is that portion of a selectin polypeptide which is necessary and sufficient for binding to a selectin ligand on endothelial cells. Binding portions include extracellular fragments of selectins, optionally fused to other polypeptides, mutated variant selectin polypeptides which retain the ability to bind endothelial selectin ligands, and the like. The endothelial selectin ligand binding portions of selectins can be identified by standard methods of molecular biology and immunology. For example, extracellular fragments of a selectin polypeptide can be prepared by deletion of portions of a nucleic acid encoding the selectin polypeptide using techniques such as PCR, exonuclease digestion, restriction endonuclease digestion to prepare fragments of the nucleic acid molecules, and then cloning the extracellular fragments. Substitutions in the amino acid sequence of a selectin polypeptide to produce functionally equivalent variants of the selectin polypeptide preferably are conservative substitutions, and typically are made by alteration of a nucleic acid encoding the selectin polypeptide. Such substitutions can be made by a variety of methods known to one of ordinary skill in the art. For example, amino acid substitutions may be made by PCR-directed mutation, site-directed mutagenesis according to the method of Kunkel (*Proc. Natl. Acad. Sci. U.S.A.* 82: 488-492, 1985), or by chemical synthesis of a gene encoding a selectin polypeptide. The activity of functionally equivalent fragments of selectin polypeptides can be tested by cloning the gene encoding the altered selectin polypeptide into a bacterial or mammalian expression vector, introducing the vector into an appropriate host cell, expressing the altered selectin polypeptide, and testing for a functional capability of the selectin polypeptides as disclosed herein. Such fragments or variants can then be tested for binding to endothelial selectin ligands using a variety of assays well known in the art, including capillary flow assays (von Andrian et al., *Cell* 82:989-999, 1995), parallel plate flow chamber assays (Lawrence and Springer, *Cell* 65:859-873, 1991; Diacovo et al., *Science*, 273: 252-255, 1996), intravital microscopy (von Andrian, *Microcirculation* 3:287-300, 1996) and the like.

As used herein with respect to dendritic cells, "expand" means to increase the number of dendritic cells in a population by culturing the population of dendritic cells or progenitors

of dendritic cells. Preferably DCs are expanded by culturing *in vitro* under defined culture conditions, such as in medium containing cytokines, growth factors and other nutrients required for DC proliferation and/or maturation.

As used herein, "transfect" or "transfection" means the introduction of one or more
5 exogenous nucleic acids into a cell. Transfection includes introduction of naked nucleic acids such as plasmids by standard physical and chemical transfection techniques, including calcium phosphate precipitation, dextran sulfate precipitation, electroporation, liposome-mediated nucleic acid transfer, ballistic methods such as particle bombardment, etc. Transfection also includes introduction of nucleic acids into cells by biological methods,
10 including viral transduction or infection (receptor-mediated and non-receptor-mediated).

In certain embodiments, isolated dendritic cells are treated (i.e., contacted *in vitro*, *ex vivo*, *in vivo*) with isolated activated platelets or membrane microparticles thereof under conditions which permit binding of the activated platelets or microparticles to the dendritic cells. Activated platelets express cell surface P-selectin, which binds to the cell-surface mucin
15 P-selectin glycoprotein ligand-1 (PSGL-1) of dendritic cells. Methods for activating platelets are well known in the art, including contacting isolated platelets with thrombin or thrombin receptor activating protein (TRAP). Membrane fractions of activated platelets, such as membrane microparticles, which include cell surface P-selectin also can be used for treating dendritic cells. The activated platelets or platelet membrane microparticles bound to the
20 dendritic cells also bind to endothelial selectin ligands present on high endothelial venules. Thus, the activated platelets or platelet membrane microparticles serve a targeting function for the isolated DCs, presumably in the same way that DC cell surface selectin expression does. In addition to the activated platelets or membrane microparticles thereof, one can use other particles containing the appropriate binding molecules to serve as a bridge between the
25 dendritic cells and endothelial selectin ligands, such as membrane vesicles (e.g. liposomes) having surface P-selectin, L-selectin, E-selectin and/or endothelial selectin ligand binding chimeras thereof. The requirements for such targeting particles are that they include (1) molecules that bind dendritic cells (e.g., P-selectin binding of PSGL-1 on DC), and (2) molecules that bind a selectin ligand on HEV (e.g., L-selectin, E-selectin, P-selectin, E-selectin/L-selectin chimeras or P-selectin/L-selectin chimeras).
30

In certain embodiments, isolated dendritic cells are loaded with one or more antigens. An antigen, as used herein, includes any molecule presented to T cells by dendritic cells,

which preferably are antigenic peptides, lipids, carbohydrates, glycolipids, or other small biological molecules. As used herein, "loaded" or "loading" means contacting dendritic cells, preferably *ex vivo*, with antigen for presentation by MHC molecules or CDI molecules. In some embodiments, the dendritic cells can be treated prior to loading to remove bound peptides from MHC molecules, e.g., existing cell-surface MHC class I molecules. In other embodiments, antigen is contacted with the dendritic cells for uptake by the DCs, degradation and presentation by MHC molecules. In all embodiments, dendritic cells that are loaded with antigen display on the cell surface complexes of MHC molecules or CDI molecules and antigen for presentation to T cells.

Presentation of certain antigens by dendritic cells also can be induced by transfecting the DCs with nucleic acid molecules which encode one or more peptide antigens. The encoded antigen may be an entire protein, of which only a small portion is antigenic (e.g., an antigenic peptide), or the antigen may be a fragment of a protein, such as an antigenic peptide. Polypeptides which include more than an antigenic peptide will be processed by the DC to the appropriate size for presentation by MHC molecules.

Antigens which are loaded as polypeptides or transfected as nucleic acid can include more than one antigen linked together (polytope), as described in greater detail below.

As used herein, "isolated" means separated from its native environment and present in sufficient quantity to permit its identification or use. As used herein with respect to cells, isolated means removed from other leukocytes present in a tissue such as blood, bone marrow, etc. As used herein, "cultured" dendritic cells are cells expanded *in vitro* from precursor cells. Isolated and/or cultured DCs preferably are substantially pure, but need not be for the methods and compositions of the invention. Substantially pure populations of dendritic cells can be prepared by techniques well known in the art including immunoaffinity purification using chromatography or magnetic separation schemes, gradient density centrifugation and the like. Dendritic cells which are stimulated to differentiate from a purified population of precursor cells also are considered substantially pure.

As used herein with respect to nucleic acids, the term "isolated" means: (i) amplified *in vitro* by, for example, polymerase chain reaction (PCR); (ii) recombinantly produced by cloning; (iii) purified, as by cleavage and gel separation; or (iv) synthesized by, for example, chemical synthesis. An isolated nucleic acid is one which is readily manipulable by recombinant DNA techniques well known in the art. Thus, a nucleotide sequence contained

in a vector in which 5' and 3' restriction sites are known or for which polymerase chain reaction (PCR) primer sequences have been disclosed is considered isolated but a nucleic acid sequence existing in its native state in its natural host is not. An isolated nucleic acid may be substantially purified, but need not be. For example, a nucleic acid that is isolated within a cloning or expression vector is not pure in that it may comprise only a tiny percentage of the material in the cell in which it resides. Such a nucleic acid is isolated, however, as the term is used herein because it is readily manipulable by standard techniques known to those of ordinary skill in the art. An isolated nucleic acid as used herein is not a naturally occurring chromosome.

As used herein, isolated, when referring to a protein or polypeptide, means, for example: (i) selectively produced by expression cloning or (ii) purified as by chromatography or electrophoresis. Isolated proteins or polypeptides may, but need not be, substantially pure. The term "substantially pure" means that the proteins or polypeptides are essentially free of other substances with which they may be found in nature or *in vivo* systems to an extent practical and appropriate for their intended use. Substantially pure polypeptides may be produced by techniques well known in the art. Because an isolated protein may be admixed with a pharmaceutically acceptable carrier in a pharmaceutical preparation, the protein may comprise only a small percentage by weight of the preparation. The protein is nonetheless isolated in that it has been separated from the substances with which it may be associated in living systems, i.e. isolated from other proteins.

In general, *ex vivo* immunotherapy involves the introduction *in vitro* of a nucleic acid which encodes an antigen into a cell(s) of a subject, and returning the genetically engineered cell(s) to the subject to stimulate an immune response. In *ex vivo* immunotherapy using dendritic cells, dendritic cell precursors are removed from a subject for *in vitro* differentiation and expansion. Nucleic acids encoding endothelial selectin ligand binding polypeptides, antigens or antigen precursors, cytokines and/or detectable markers are introduced (i.e., transduced or transfected) into the cells *in vitro*. Typically, the modified cells are then expanded in culture before being reimplanted into a subject. As used herein, "subject" means a mammal, including a human, a non-human primate, a sheep, a goat, a horse, a cow, a dog, a cat, and a rodent. The preferred subject is a human.

The compositions and methods of the invention are useful for modulating the responsiveness of a subject's immune system to one or more antigens, e.g. by inducing

increased production of cytotoxic T lymphocytes which recognize the antigens or by inducing tolerance of the antigens. This is useful for modulating a subject's immune response for the treatment of tumors or pathogenic infections (increased immune response), or for treatment of autoimmune diseases, allergy, or transplantation (decreased immune response).

The invention is not limited in utility to human immunotherapy, but also provides a method for assessing the effects of immunosuppressive agents in mammalian models such as primates, pigs, sheep, dogs, rodents, and cows. The invention also provides an improved method for testing the effectiveness of antigenic peptides in a mammal. A series of peptides can be administered to the group of mammals as immunizations. The series of peptides can include variants of recognized antigenic peptides and the immune response of each of these variants can thus be assessed to determine the optimal amino acid sequence of immune response boosting peptides. Modifications to the peptides can be made based on known parameters of HLA binding affinity described herein, or randomly, and tested for immunogenic potential by the same methodology. *In vitro* uses also will be known to one of ordinary skill in the art, including presentation of antigen to T cells for *in vitro* maturation and stimulation protocols.

Exemplary tumor associated peptide antigens that can be expressed to induce or enhance an immune response are derived from tumor associated genes and encoded proteins including MAGE proteins, MART, LAGE, NY-ESO-1, tyrosinase, PRAME, prostate specific antigen (PSA), BCR-ABL, immunoglobulins from B cell tumors, T cell receptors from T cell tumors, RAS oncogene products, carcinoembryonic antigen (CEA), mutated p53, Melan-A, tyrosinase-related protein-1 and the like.

Other examples will be known to one of ordinary skill in the art (for example, see Coulie, *Stem Cells* 13:393-403, 1995), and can be used in the invention in a like manner as those disclosed herein.

Antigens that are characteristic of pathogens include antigens derived from viruses, bacteria, parasites or fungi. Examples of important pathogens include *Vibrio cholerae*, enterotoxigenic *Escherichia coli*, rotavirus, *Clostridium difficile*, *Shigella* species, *Salmonella typhi*, parainfluenza virus, influenza virus, *Streptococcus pneumoniae*, *Borella burgdorferi*, HIV, *Streptococcus mutans*, *Plasmodium falciparum*, *Staphylococcus aureus*, rabies virus and Epstein-Barr virus.

Viruses in general include but are not limited to those in the following families:

picornaviridae; caliciviridae; togaviridae; flaviviridae; coronaviridae; rhabdoviridae; filoviridae; paramyxoviridae; orthomyxoviridae; bunyaviridae; arenaviridae; reoviridae; retroviridae; hepadnaviridae; parvoviridae; papovaviridae; adenoviridae; herpesviridae; and poxyviridae.

- 5 Bacteria in general include but are not limited to: *P. aeruginosa*; *E. coli*; *Klebsiella* sp.; *Serratia* sp.; *Pseudomonas* sp.; *P. cepacia*; *Acinetobacter* sp.; *S. epidermis*; *E. faecalis*; *S. pneumoniae*; *S. aureus*; *Haemophilus* sp.; *Neisseria* sp.; *N. meningitidis*; *Bacteroides* sp.; *Citrobacter* sp.; *Branhamella* sp.; *Salmonella* sp.; *Shigella* sp.; *S. pyogenes*; *Proteus* sp.; *Clostridium* sp.; *Erysipelothrix* sp.; *Lesteria* sp.; *Pasteurella multocida*; *Streptobacillus* sp.;
10 *Spirillum* sp.; *Fusospirocheta* sp.; *Treponema pallidum*; *Borrelia* sp.; *Actinomycetes*; *Mycoplasma* sp.; *Chlamydia* sp.; *Rickettsia* sp.; *Spirochaeta*; *Legionella* sp.; *Mycobacteria* sp.; *Ureaplasma* sp.; *Streptomyces* sp.; *Trichomoras* sp.; and *P. mirabilis*.

- Parasites include but are not limited to: *Plasmodium falciparum*, *P. vivax*, *P. ovale*, *P. malaria*; *Toxoplasma gondii*; *Leishmania mexicana*, *L. tropica*, *L. major*, *L. aethiopica*, *L. donovani*; *Trypanosoma cruzi*, *T. brucei*; *Schistosoma mansoni*, *S. haematobium*, *S. japonium*;
15 *Trichinella spiralis*; *Wuchereria bancrofti*; *Brugia malayi*; *Entamoeba histolytica*; *Enterobius vermicularis*; *Taenia solium*, *T. saginata*; *Trichomonas vaginalis*, *T. hominis*, *T. tenax*; *Giardia lamblia*; *Cryptosporidium parvum*; *Pneumocystis carinii*; *Babesia bovis*, *B. divergens*, *B. microti*; *Isospora belli*, *I. hominis*; *Dientamoeba fragilis*; *Onchocerca volvulus*; *Ascaris lumbricoides*; *Necator americanus*; *Ancylostoma duodenale*; *Strongyloides stercoralis*;
20 *Capillaria philippinensis*; *Angiostrongylus cantonensis*; *Hymenolepis nana*; *Diphyllobothrium latum*; *Echinococcus granulosus*, *E. multilocularis*; *Paragonimus westermani*, *P. caliensis*; *Chlonorchis sinensis*; *Opisthorchis felinus*, *O. viverrini*; *Fasciola hepatica*; *Sarcoptes scabiei*; *Pediculus humanus*; *Phthirus pubis*; and *Dermatobia hominis*.

- 25 Fungi in general include but are not limited to: *Cryptococcus neoformans*; *Blastomyces dermatitidis*; *Ajellomyces dermatitidis*; *Histoplasma capsulatum*; *Coccidioides immitis*; *Candida* species, including *C. albicans*, *C. tropicalis*, *C. parapsilosis*, *C. guilliermondii* and *C. krusei*; *Aspergillus* species, including *A. fumigatus*, *A. flavus* and *A. niger*; *Rhizopus* species; *Rhizomucor* species; *Cunninghamella* species; *Apophysomyces* species, including *A. saksenaeae*, *A. mucor* and *A. absidia*; *Sporothrix schenckii*;
30 *Paracoccidioides brasiliensis*; *Pseudallescheria boydii*; *Torulopsis glabrata*; and *Dermatophytes* species.

Specific examples of antigens characteristic of a pathogen include: the influenza virus nucleoprotein (residues 218-226, Fu et al., *J. Virol.* 71:2715-2721, 1997), antigens from Sendai virus and lymphocytic choriomeningitis virus (An et al., *J. Virol.* 71:2292-2302, 1997), the E1 protein of hepatitis C virus (Bruna-Romero et al., *Hepatology* 25:470-477, 1997), the virus envelope glycoprotein gp160 of HIV (Achour et al., *J. Virol.* 70: 6741-6750, 1996), amino acids 252-260 or the circumsporozoite protein of *Plasmodium berghei* (Allsopp et al., *Eur. J. Immunol.* 26:1951-1958, 1996), the influenza A virus nucleoprotein (residues 366-374, Nomura et al., *J. Immunol. Methods* 193:4149, 1996), the listeriolysin O protein of *Listeria monocytogenes* (residues 91-99, An et al., *Infect. Immun.* 64:1685-1693, 1996), the E6 protein (residues 131-140, Gao et al., *J. Immunol.* 155:5519-5526, 1995) and E7 protein (residues 21-28 and 48-55, Bauer et al., *Scand. J. Immunol.* 42:317-323, 1995) of human papillomavirus type 16, the M2 protein of respiratory syncytial virus (residues 82-90 and 81-95, Hsu et al., *Immunology* 85:347-350, 1995), the herpes simplex virus type 1 ribonucleotide reductase (Salvucci, et al., *J. Virol.* 69:1122-1131, 1995) and the rotavirus VP7 protein (Franco et al., *J. Gen. Virol.* 74:2579-2586, 1993).

Administration of antigens via modified and/or treated dendritic cells with certain cytokines which induce tolerance (e.g., interleukin-10, TGF- β) is useful for reducing immune responses to those antigens. This is useful in the treatment of allergic responses and autoimmune diseases, as well as in treating allograft rejection.

Antigens that are allergens are generally proteins or glycoproteins, although allergens may also be low molecular weight allergenic haptens that induce allergy after covalently combining with a protein carrier (Remington's Pharmaceutical Sciences). Allergens include antigens derived from pollens, dust, molds, spores, dander, insects and foods. Specific examples include: the major horse allergen Equ c1 (Gregoire et al., *J. Biol. Chem.* 271:32951-32959, 1996), the Hor v 9 pollen allergen from barley (Astwood et al., *Gene* 182:53-62, 1996), the major allergen of the domestic cat, Fel d 1 (Counsell et al., *J. Allergy Clin. Immunol.* 98:884-894, 1996), a major latex allergen, Hev b 5 (Salter et al., *J. Biol. Chem.* 271:25394-25399, 1996), a major allergen of salmon, Sal s 1 (Lindstrom et al., *Scand. J. Immunol.* 44:335-344, 1996), allergens of the house dust mite, *B. tropicalis* (Carabello et al., *J. Allergy Clin. Immunol.* 98:573-579, 1996; Der p 2, Chua et al., *Clin. Exp. Allergy* 26:829-837, 1996; Der p 2, Harris et al., *Int. Immunol.* 9:273-280, 1997), major allergens of velvet grass, Hol 1 1 and Hol 1 5 (Schramm et al., *Int. Arch. Allergy immunol.* 110:354-363, 1996), a

major allergen of ryegrass, Lol p 9 (Blaher et al., *J. Allergy Clin. Immunol.* 98:124-132, 1996), the predominant allergen of bovine dander, BDA20 (Mantyljarvi et al., *J. Allergy Clin. Immunol.* 97:1297-1303, 1996), a major allergen of Kentucky bluegrass pollen, rKBG60 (Zhang et al., *Immunology*, 87:283-290, 1996), and the hornet venom allergen Dol m 5 (King et al., *J. Allergy Clin. Immunol.* 99:630-639, 1997).

Examples of antigens characteristic of autoimmune disease include: antigens from human myelin basic protein (residues 110-118), proteolipid protein (residues 80-88), myelin-associated protein (residues 287-295, 509-517, 556-564) in multiple sclerosis (Tsuchida et al. *Proc. Natl. Acad. Sci USA* 91:10859-10863, 1994), the islet cell antigen ICA69 in diabetes (Karges et al., *Biochim. Biophys. Acta* 1360:97-101, 1997), the protein L7 in rheumatic diseases such as systemic lupus erythematosus, rheumatoid arthritis, and systemic sclerosis (Neu et al., *Arthritis Rheum.* 40:661-671, 1997), two peptides from human acetylcholine receptor alpha-subunit, p195-212 and p259-271 in myasthenia gravis (Katz-Levy et al., *Proc. Natl. Acad. Sci. U S A* 94: 3200-3205, 1997), the nuclear autoantigen La (SS-B) protein (Bachmann et al., *J. Autoimmun.* 9:747-756, 1996) and the 52-kd Ro(SS-A) protein (Dorner et al., *Hepatology* 24:1404-1407, 1996) in Sjogren's syndrome, and cytochrome P450IID6, the main target antigen of LKM-1 antibody-positive type II autoimmune hepatitis (Lohr et al., *Hepatology* 24:1416-1421, 1996). Other autoimmune antigens will be known to one of ordinary skill in the art.

Other peptide antigens can be identified according to methods used in the foregoing references which describe antigens of tumors, pathogens, allergens and autoimmune disease. Standard references in the art (e.g., *Remington's Pharmaceutical Sciences*, 18th edition, 1990) provide modes of administration and formulations for delivery of immunogens with adjuvant or in a non-adjuvant carrier.

As noted previously, antigens can be provided as polypeptides, peptides (i.e. in a form and length which can be bound and presented by MHC molecules), nucleic acid molecules which encode peptide antigens, lipids, carbohydrates, glycolipids, or other small biological molecules. Especially preferred are nucleic acids encoding a series of antigenic epitopes, known as "polytopes". The epitopes can be arranged in sequential or overlapping fashion (see, e.g., Thomson et al., *Proc. Natl. Acad. Sci. USA* 92:5845-5849, 1995; Gilbert et al., *Nature Biotechnol.* 15:1280-1284, 1997), with or without the natural flanking sequences, and can be separated by unrelated linker sequences if desired. The polytope is processed to

generated individual epitopes which are recognized by the immune system for generation of immune responses.

Thus, for example, several HLA class I and HLA class II antigens can be combined (e.g. by preparation of hybrid nucleic acids or polypeptides) to form "polytopes", according to standard procedures of molecular biology. Polytopes comprising two or more potentially immunogenic or immune response stimulating peptides can be joined together in various arrangements (e.g. concatenated, overlapping). The polytope (or nucleic acid encoding the polytope) can be administered in vaccination protocols using the dendritic cells disclosed herein, e.g. to animals, to test the effectiveness of the polytope in stimulating, enhancing and/or provoking an immune response.

The peptides can be joined together directly or via the use of flanking sequences to form polytopes, and the use of polytopes as vaccines is well known in the art (see, e.g., Thomson et al., *Proc. Acad. Natl. Acad. Sci USA* 92(13):5845-5849, 1995; Gilbert et al., *Nature Biotechnol.* 15(12):1280-1284, 1997; Thomson et al., *J. Immunol.* 157(2):822-826, 1996; Tam et al., *J. Exp. Med.* 171(1):299-306, 1990). For example, Tam showed that polytopes consisting of both MHC class I and class II binding epitopes successfully generated antibody and protective immunity in a mouse model. Tam also demonstrated that polytopes comprising "strings" of epitopes are processed to yield individual epitopes which are presented by MHC molecules and recognized by CTLs. Thus polytopes containing various numbers and combinations of epitopes can be prepared and tested for recognition by CTLs and for efficacy in increasing an immune response.

It is known, for example, that tumors express a set of tumor antigens, of which only certain subsets may be expressed in the tumor of any given patient. Polytopes can be prepared which correspond to the different combination of epitopes representing the subset of tumor rejection antigens expressed in a particular patient. Polytopes also can be prepared to reflect a broader spectrum of tumor rejection antigens known to be expressed by a tumor type. Likewise, combinations of pathogen antigens can be made to induce protective immune responses against a plurality of pathogens.

Polytopes can be introduced to a patient in need of such treatment as polypeptide structures, or via the use of nucleic acid delivery systems known in the art (see, e.g., Allsopp et al., *Eur. J. Immunol.* 26(8):1951-1959, 1996). Adenovirus, retroviruses, adeno-associated virus, plasmids, bacteria, etc. can be used in such delivery. One can test the polytope delivery

systems in mouse models to determine efficacy of the delivery system. The systems also can be tested in human clinical trials.

As part of the immunization protocols, substances which modulate the immune response may be administered with nucleic acid or peptide components of a vaccine. Such immune response modulating compounds are referred to herein as "adjuvants". Adjuvants of many kinds are well known in the art; specific examples include MPL, a congener obtained after purification and acid hydrolysis of *Salmonella minnesota* Re 595 lipopolysaccharide, saponins including QS21, a pure QA-21 saponin purified from *Quillja saponaria* extract, DQS21, described in PCT application WO96/33739 (SmithKline Beecham), vitamin E and various water-in-oil emulsions prepared from biodegradable oils such as squalene and/or tocopherol.

Cytokines are also useful in vaccination protocols as adjuvants. Many cytokines useful as a result of their lymphocyte regulatory properties will be known to one of ordinary skill in the art, including interleukin-12 (IL-12) which has been shown to enhance the protective effects of vaccines (*Science* 268: 1432-1434, 1995), GM-CSF and IL-18. Adjuvants which inhibit the immune response are useful in vaccination protocols for tolerization to antigens. Such adjuvants include factors such as IL-10 (see, e.g., Steinbrink et al., *J. Immunol.* 159:4772-80, 1997; Takayama et al., *Transplantation* 66:1567-74, 1998) and TGF- β .

There are a number of additional immune response potentiating compounds that can be used in vaccination protocols. These include costimulatory molecules provided in either protein or nucleic acid form. Such costimulatory molecules include the B7-1 and B7-2 (CD80 and CD86 respectively) molecules which are expressed on dendritic cells and interact with the CD28 molecule expressed on the T cell. This interaction provides costimulation (signal 2) to an antigen/MHC/TCR stimulated (signal 1) T cell, increasing T cell proliferation and effector function. B7 also interacts with CTLA4 (CD152) on T cells and studies involving CTLA4 and B7 ligands indicate that the B7-CTLA4 interaction can enhance antitumor immunity and CTL proliferation (Zheng et al., *Proc. Natl. Acad. Sci. USA* 95:6284-6289, 1998). In addition, a combination of B7/IL-6/IL-12 costimulation has been shown to induce IFN-gamma and a Th1 cytokine profile in the T cell population leading to further enhanced T cell activity (Gajewski et al., *J. Immunol.* 154:5637-5648, 1995). The use of anti-CD28 antibodies to directly stimulate T cells could also be considered.

Lymphocyte function associated antigen-3 (LFA-3) is expressed on APCs and some tumor cells and interacts with CD2 expressed on T cells. This interaction induces T cell IL-2 and IFN-gamma production and can thus complement but not substitute, the B7/CD28 costimulatory interaction (Parra et al., *J. Immunol.*, 158:637-642, 1997; Fenton et al., *J. Immunother.*, 21:95-108, 1998).

Lymphocyte function associated antigen-1 (LFA-1) is expressed on leukocytes and interacts with ICAM-1 (CD54) expressed on APCs. This interaction induces T cell IL-2 and IFN-gamma production and can thus complement but not substitute, the B7/CD28 costimulatory interaction (Fenton et al., 1998). ICAM-1 is thus a further example of a costimulatory molecule that could be provided in a vaccination protocol in the various ways discussed above for B7.

Still other examples of molecules that could be provided to dendritic cells in a vaccination protocol are chemokine receptors including CC chemokine receptor 7 (CCR7), which is a receptor for secondary lymphoid-tissue chemokine (SLC) and EBI1 ligand chemokine (ELC).

Complete CTL activation and effector function requires Th cell help through the interaction between the Th cell CD40L (CD40 ligand) molecule and the CD40 molecule expressed by DCs (Ridge et al., *Nature* 393:474, 1998; Bennett et al., *Nature* 393:478, 1998; Schoenberger et al., *Nature* 393:480, 1998). This mechanism of this costimulatory signal is likely to involve upregulation of B7 and associated IL-6/IL-12 production by the DC (APC). The CD40-CD40L interaction thus complements the signal 1 (antigen/MHC-TCR) and signal 2 (B7-CD28) interactions.

The cultured dendritic cells described herein can be assayed for expression of these molecules, and supplemented if necessary by, e.g., transfection of appropriate nucleic acids, such as by nucleic acid (naked DNA) immunization (Kim et al., *Nature Biotechnol.* 15:7:641-646, 1997) or recombinant viruses (Wendtner et al., *Gene Ther.* 4:726-735, 1997). These systems are all amenable to the construction and use of expression cassettes for the coexpression of B7 or other stimulatory molecules with other molecules of choice such as the antigens or fragment(s) of antigens discussed herein (including polypeptides) or cytokines. These delivery systems can be used for induction of the appropriate molecules *in vitro* and for *in vivo* vaccination situations.

The use of anti-CD40 antibodies to stimulate DCs directly, would be expected to

enhance a response to antigens which are normally encountered outside of an inflammatory context. Other methods for inducing maturation of dendritic cells, e.g., by increasing CD40-CD40L interaction, or by contacting DCs with CpG-containing oligodeoxynucleotides or stimulatory sugar moieties from extracellular matrix, are known in the art. In these situations Th help and B7 costimulation signals are not provided. This mechanism might be used in the context of antigen loaded DC based therapies or in situations where Th epitopes have not been defined within known antigen precursors.

As used herein, antigen embraces a naturally selected antigenic molecule (peptide, glycolipid, other small molecules, etc.) presented by MHC or CD1 molecules, as well as variants of the naturally selected antigenic molecules which can be prepared according to methods known in the art and disclosed herein. For example, peptide antigens administered according to the methods described herein can have the amino acid sequence of a naturally processed antigenic peptide. Alternatively, modifications can be made to the amino acid sequence to enhance binding to HLA for presentation to a subject's immune system, while retaining the ability to induce CTLs which recognize the naturally processed peptide. The portions of HLA binding peptides important for binding activity are known (*see, e.g., Parker et al., J. Immunol.* 149:3580-3587, 1992). Variant antigenic peptides which bind more tightly to HLA molecules have been found to be more immunogenic (Parkhurst et al., *J. Immunol.* 157:2539-2548, 1996; Bakker et al., *Int. J. Cancer* 70:302-309, 1997). A general protocol for modification of HLA binding peptides has been suggested by Parker et al. (*J. Immunol.* 152:163-175, 1994). Thus one of ordinary skill in the art, with only routine experimentation, can design synthetic variant peptides which can be administered to induce an immune response in a host. The methods disclosed herein provide a sensitive assay for determining the immunogenic potential of such modified peptides.

The peptides used in accordance with the invention can be prepared by any method known in the art. Typically this entails programming a peptide synthesizer with a desired amino acid sequence, providing the subunits to be incorporated into the extending peptide chain, and purifying the synthesized peptide. All of these steps are well known and practiced routinely in the art. The peptides thus synthesized can incorporate modified amino acids and/or modified inter-amino acid bonds, if desired, to increase stability, reduce proteolysis, or confer some other property to the peptides. Of course, peptides may be made by other methods, including transcription and translation of a nucleic acid which encodes such

peptides.

Antigens may also be expressed by the dendritic cells. For example, dendritic cells can be transfected with a nucleic acid molecule encoding the antigen or the polypeptide from which the antigen is derived. The use of a complete protein in this manner may be preferable because the host alleles can select the relevant epitope or epitopes for presentation. Host epitope selection recently was demonstrated by Zhai et al (*J. Immunol.* 156: 700-710, 1996), wherein tumor infiltrating lymphocytes recognized five different epitopes presented on the surface of cells transduced with a viral vector expressing a cancer antigen. Alternatively, the use of only a portion of a gene which does not encode a functional protein, e.g., an antigen or antigen precursor, may be preferable where it is suspected that the protein may have deleterious effects. Expressing only a portion of the gene permits vaccination against an antigen without concomitant expression of a complete functional gene product in the dendritic cells.

In preferred embodiments, the selectin polypeptide is introduced (e.g., transfected, transduced) into dendritic cells by a viral vector selected from the group consisting of adenoviruses, retroviruses, adeno-associated viruses, poxviruses including vaccinia viruses and attenuated poxviruses, lentiviruses including HIV and HIV-derived viruses, Semliki Forest virus, Venezuelan equine encephalitis virus, Sindbis virus, lambda bacteriophage and Ty virus-like particle. Examples of viruses and virus-like particles which have been used to deliver exogenous nucleic acids include: replication-defective adenoviruses (e.g., Xiang et al., *Virology* 219:220-227, 1996; Eloit et al., *J. Virol* 71:5375-5381, 1997; Chengalvala et al., *Vaccine* 15:335-339, 1997), a modified retrovirus (Townsend et al., *J. Virol.* 71:3365-3374, 1997), a nonreplicating retrovirus (Irwin et al., *J. Virol.* 68:5036-5044, 1994), a replication defective Semliki Forest virus (Zhao et al., *Proc. Natl. Acad. Sci. USA* 92:3009-3013, 1995), canarypox virus and highly attenuated vaccinia virus derivative (Paoletti, *Proc. Natl. Acad. Sci. USA* 93:11349-11353, 1996), non-replicative vaccinia virus (Moss, *Proc. Natl. Acad. Sci. USA* 93:11341-11348, 1996), replicative vaccinia virus (Moss, *Dev. Biol. Stand.* 82:55-63, 1994), Venezuelan equine encephalitis virus (Davis et al., *J. Virol.* 70:3781-3787, 1996), Sindbis virus (Pugachev et al., *Virology* 212:587-594, 1995) and modified bacteriophage lambda (PCT/US97/12928, WO96/21007). Antigens and/or cytokines also may be expressed in dendritic cells using such vectors.

Preferred viral vectors are replication defective. As used herein, a "replication-

defective" virus or viral vector is one which is incapable of replicating autonomously in the target cell. Generally, the genome of a replication-defective adenovirus used in the context of the present invention contains mutations or deletions of at least the sequences needed for replication of the adenovirus in the infected cell. Such sequences are well known to those of ordinary skill in the art, and include, for example, in adenoviruses portions of the E1, E3, and E4 regions of the genome.

In one preferred embodiment, the virus vector is an adenovirus. An "adenovirus", for the delivery of nucleic acids encoding selectin polypeptides, refers to an adenovirus that: (1) contains exogenous genetic material that can be transcribed and translated in a mammalian cell and which encodes a polypeptide that binds a selectin ligand on endothelial cells, i.e., a selectin polypeptide, and (2) contains on its surface a ligand that selectively binds to a receptor on the surface of a target dendritic cell, and thereby gains entry to the target cell. The term adenovirus also embraces an adenovirus genome containing exogenous genetic material which encodes a polypeptide that binds a selectin ligand on endothelial cells, i.e. an unencapsidated adenovirus. As used herein, "exogenous genetic material" refers to a nucleic acid molecule (e.g., nucleic acid or oligonucleotide), either natural or synthetic, that is not naturally found in an adenovirus. The "exogenous genetic material" includes a gene or fragment thereof which encodes a polypeptide that binds a selectin ligand on endothelial cells, an antigen or precursor thereof that can, if necessary, be processed into one or more antigens, or a cytokine.

The complete nucleotide sequences of adenovirus genomes are known and have been deposited in nucleotide sequence databases. For example, the genome of the adenovirus type 5 has been completely sequenced and is accessible via GenBank accession number M73260. Similarly, portions or even whole genomes of other adenovirus types (type 2, type 7, type 12, and the like), retroviruses, and other viral vectors have also been sequenced and deposited in databases.

The nucleic acid encoding an antigen or precursor thereof preferably is inserted into a region of the virus genome which is not essential to the production of replication-defective recombinant viruses. For example, the nucleic acid preferably is not inserted into regions which contain adenovirus genes encoding proteins which are not easily supplied *in trans*. Thus, the nucleic acid preferably is inserted into the E1 region, which can be complemented (supplied *in trans*) by an adenovirus encapsidation cell line such as 293 cells. Other preferred

5 sites of insertion of the nucleic acid include the E3 region, which is not required for production of replication-defective recombinant adenoviruses, and the E4 region, mutation of which can be complemented by co-transduction with a helper virus or plasmid or by infection of a suitable complementary cell line. Other sites also may be used as will be apparent to one of ordinary skill in the art. In particular, access to the nucleotide sequences of virus genomes enables a person skilled in the art to identify regions of the virus genomes suitable for insertion of the nucleic acid encoding a selectin polypeptide, and/or antigen or antigen precursor, and/or a cytokine.

10 The nucleic acids assembled to prepare a complete replication-defective virus genome or other viral or non-viral vector can be prepared by any method known in the art. For example, a virus genome or plasmid can be isolated and then modified *in vitro* by standard methods of molecular biology (see, e.g., *Molecular Cloning: A Laboratory Manual*, J. Sambrook, et al., eds., Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989, or *Current Protocols in Molecular Biology*, F.M. Ausubel, et al., eds., John Wiley & Sons, Inc., New York). The modified virus genome so obtained optionally can be isolated and used to transfect an encapsidation cell line if necessary.

15 In the preferred embodiments, the virus genome further includes a regulatory sequence, e.g., a promoter region (also referred to as a "promoter"), that is operably coupled to the nucleic acid molecule encoding an antigen or antigen precursor. The regulatory sequence controls the expression of the nucleic acid molecule encoding a selectin polypeptide, and/or an antigen or antigen precursor, and/or a cytokine in the target dendritic cell. As used herein, a nucleic acid molecule encoding one or more polypeptides (the "coding sequence") and regulatory sequences are said to be "operably" joined when they are covalently linked in such a way as to place the transcription or the expression of the coding sequence under the influence or control of the regulatory sequences. If it is desired that the coding sequence be translated into a functional protein, two DNA sequences are said to be operably joined if induction of a promoter in the 5' regulatory sequence results in the transcription of the coding sequence and if the nature of the linkage between the two DNA sequences does not (1) result in the introduction of a frame-shift mutation, (2) interfere with the ability of the promoter region to direct the transcription of the coding sequences, or (3) interfere with the ability of the corresponding RNA transcript to be translated into a protein. Thus, a promoter region would be operably joined to a coding sequence if the promoter region were capable of

effecting transcription of that DNA sequence such that the resulting transcript might be translated into the desired protein or polypeptide.

The precise nature of the regulatory sequences needed for gene expression may vary between species or cell types, but shall in general include, as necessary, 3' or 5' non-transcribed and non-translated sequences involved with the initiation of transcription and translation respectively, such as a TATA box, CAAT sequence, and the like. In particular, such 5' non-transcribed regulatory sequences will include a promoter region which includes a promoter sequence for transcriptional control of the operably joined gene. Regulatory sequences can also include enhancer sequences or upstream 5' or downstream 3' transcriptional regulatory sequences as desired.

Exemplary promoters that are useful in the invention include constitutive promoters and regulatable promoters (e.g., cell lineage specific promoters, inducible promoters). Exemplary constitutive promoters include promoters derived from cytomegalovirus, a long terminal repeat (LTR) of retroviruses, e.g., Rous sarcoma virus or Moloney murine leukemia virus, and adenovirus E1A promoter, an adenovirus MLP promoter and a SR α promoter. Exemplary tissue or cell specific transcriptional regulatory sequences are those which are active in dendritic cells, including CD11a, dectins-1 and 2, MHC class II, CD1a, b or c, CD80 and CD86. Exemplary inducible promoters are described in the following references: *Science* 268:1786 (1995); *TIBS* 18:471 (1993); *PNAS* 91:3180 (1994); *PNAS* 90:1657 (1993); *PNAS* 88:698 (1991); *Nature Biotechnol.* 14:486 (1996); and *PNAS* 93:5185 (1996). An exemplary repressible promoter, the tetracycline repressible system, is described in *PNAS* 89:5547 (1992). Other constitutive, tissue-specific, inducible and repressible promoters will be known by those of skill in the art and thus are not listed here.

The expression vectors (plasmids, viruses, etc.) optionally contain one or more sequences that are suitable for use in the identification of cells that have or have not been transfected or transduced. "Transfection", as used herein, refers to the introduction of a plasmid or other non-viral nucleic acid molecule into the target cell. "Transduction", as used herein, refers to the introduction of the virus genome into the target cell. Markers to identify cells that have been transfected or transduced include, for example, genes encoding proteins that increase or decrease resistance or sensitivity to antibiotics or other compounds, genes which encode enzymes having activities that are detectable by standard assays known in the art and genes which detectably (e.g. visibly) affect the phenotype of the transduced target

cells. Exemplary genes that are suitable as markers include a *lacZ* gene, a chloramphenicol acetyltransferase gene, an alkaline phosphatase gene, a luciferase gene, and a green fluorescent protein gene. Preferred markers are those which can be used as a basis for selection by fluorescence activated cell sorting or magnetic sorting.

5 Methods for delivering whole encapsidated virus include contacting dendritic cells with the virus, whereby the virus genome can be delivered by receptor-mediated endocytosis via binding of a viral capsid protein to a cellular receptor. Methods for delivering non-encapsidated viral genomes and other nucleic acid molecules such as plasmids include the foregoing methods and also methods for delivery of nucleic acids to cells familiar to those of
10 skill in the art of molecular biology. For example, when delivering a recombinant viral genome without any associated coat protein, or an expression plasmid, the nucleic acid can be introduced into a cell by transfection using a standard technique such as electroporation, liposome transfection, calcium phosphate precipitation, or a commercially available technology such as the Tfx-50 transfection reagent (Promega Corp., Madison, WI).

15 The genetically modified or platelet modified dendritic cells of the invention can be delivered to a subject by methods known to those of ordinary skill in the art, particularly parenteral administration methods. Preferably a subject is injected intraarterially or intravenously with the modified or treated dendritic cells, or with dendritic cells that have been modified genetically and with platelets. The invention also includes administration of
20 acitivated platelets to a subject prior to (preferably immediately prior to) or concurrently with genetically modified dendritic cells.

The invention provides other compositions and kits which are useful for practicing the above-described methods. According to another aspect of the invention, kits are provided which contain (a) nucleic acid molecules that encodes a selectin polypeptide; and optionally
25 (b) a peptide antigen or nucleic acid encoding an antigen, and optionally, (c) a separate container of adjuvant. Instructions for the use of the nucleic acid encoding the selectin polypeptide and the antigen(s) can also be included. Other kits are like those described above except that they include instructions for preparing isolated activated platelets or membrane microparticles thereof, and treating the dendritic cells with the platelets or microparticles.
30 Still other kits are like those described above except that they include instructions for administering isolated activated platelets or membrane microparticles thereof prior to or concurrently with genetically modified dendritic cells. The components of the kits are

sufficient, when used, for example, to modify dendritic cells isolated from a subject and subsequently administered to a subject, to modulate an immune response in the subject against an antigen presented by the modified dendritic cells.

Methods for increasing or decreasing immune responses to antigens are also provided.

5 The methods utilize the unique features of dendritic cells as antigen-presenting cells, and in particular, of modified isolated cultured dendritic cells that either express a selectin polypeptide sufficient to permit targeting to peripheral lymph nodes, and/or which have been modified by binding of activated platelets or microparticles (*in vitro*, *ex vivo*, or *in vivo*). These features permit enhanced presentation of antigens to T lymphocytes in lymph nodes, 10 which under certain conditions will increase an immune response to the antigens, and under other conditions will decrease an immune response to the antigens. Thus the methods contemplate loading modified dendritic cells with antigen and using such cells for efficient delivery of antigen to the immune system.

15 The pharmaceutical compositions used in the methods should be sterile and contain a therapeutically effective amount of the modified dendritic cells and optionally activated platelets or microparticles thereof for modulating an immune response in a unit of weight or volume suitable for administration to a patient. In general, an effective amount is that amount of a pharmaceutical preparation that alone, or together with further doses, stimulates (or 20 decreases) the desired response. The immune response can be measured by determining the activity of cytotoxic T lymphocytes or antige-specific antibody titers after the administration of the modified dendritic cells, and preferably both before and after the administration of the dendritic cells. Methods for measuring cytotoxic T lymphocyte activity include measurement of tumor necrosis factor release by the cytotoxic T lymphocytes and measurement of 25 chromium release as is well known in the art. Other assays will be known to one of ordinary skill in the art and can be employed for measuring the level of the immune response following dendritic cell administration.

30 Where it is desired to stimulate an immune response using a therapeutic composition of the invention, this may involve the stimulation of a humoral antibody response resulting in an increase in antibody titer in serum, a clonal expansion of cytotoxic T lymphocytes, or some other desirable immunologic response. Where it is desired to inhibit an immune response using a therapeutic composition of the invention, this may involve the inhibition of a humoral antibody response resulting in a decrease in antibody titer in serum, a decrease in the number

of reactive cytotoxic T lymphocytes, or some other desirable immunologic response. The absolute amount will depend upon a variety of factors, including the material selected for administration, whether the administration is in single or multiple doses, and individual patient parameters including age, physical condition, size, weight, and the stage of the disease.

- 5 These factors are well known to those of ordinary skill in the art and can be addressed with no more than routine experimentation.

The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredients. The term "physiologically acceptable" refers to a non-toxic material that is compatible with a biological
10 system such as a cell, cell culture, tissue, or organism. The characteristics of carriers included in the compositions of the invention will depend on the route of administration.

Physiologically and pharmaceutically acceptable carriers include diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials which are well known in the art (see, e.g.,
15 *Remington's Pharmaceutical Sciences*).

The doses of modified dendritic cell and antigen administered to a subject can be chosen in accordance with different parameters, in particular in accordance with the mode of administration used and the state of the subject. Other factors include the particular antigen used for immunization and the desired period of treatment.

Examples

Example 1: Determination of Cultured Dendritic Cell Homing to Peripheral Lymph Node

Dendritic cells (DCs) are known to have a unique and potent capacity to present antigens to naive T cells in peripheral lymph nodes (PLN). Thus dendritic cells are envisioned as ideal immunotherapeutic tools for modulating immune responses. For use in
25 immunotherapy, large numbers of DCs are grown from CD34⁺ progenitors or blood monocytes by *in vitro* differentiation. Because antigen presentation to T cells occurs in the PLN, the capacity of cultured DCs to traffic to PLN was examined.

Murine dendritic cells were grown from bone marrow progenitors as described by Mayordomo et al. (*Nature Med.* 1:1297-1302, 1995) with minor modifications. Briefly, bone
30 marrow progenitors were isolated, red blood cells lysed, and the progenitors were cultured without depletion of lymphocytes in the presence of granulocyte-macrophage colony stimulating factor (GM-CSF; Immunex Corp.) and interleukin-4 (IL-4; R & D Systems) at

500 U/ml each in culture medium. For growth of human dendritic cells from cord blood CD34⁺ progenitors, the culture medium included hGM-CSF (1000U/ml), hIL-4 (50 U/ml), Flt3 ligand (100ng/ml; Immunex Corp. and R & D Systems) and TNF- α (50 U/ml; R & D Systems). For growth of human dendritic cells from human monocytes (peripheral blood), the culture medium included hGM-CSF (1000U/ml) and hIL-4 (1000 U/ml). For observation, DCs were fluorescently labeled with calcein (2.5 μ g/ml/10⁷ cells; Molecular Probes, Inc.).

The migration of the cultured dendritic cells was observed after intravenous injection. When injected intravenously, DCs did not reach the PLN, but rather accumulated in extranodal sites, mainly spleen and liver.

Example 2: Determination of L-selectin Expression on Cultured Dendritic Cells.

DC homing to PLN from blood is mediated by DC cell surface L-selectin binding to selectin ligands (peripheral node addressins, PNAds) on high endothelial venules (HEV). The expression of L-selectin and binding of the cultured DCs to PNAds were examined to determine if the defect in PLN homing of DCs was due to inefficient binding of DCs to HEV.

Dendritic cell binding to PNAds was examined in a parallel plate flow chamber assay described by Lawrence and Springer (*Cell* 65:859-873, 1991). DCs were cultured under a variety of conditions including those described above with or without several concentrations of TGF- β , IL-3 and inhibitors of metalloproteinases.

Regardless of the progenitor cell used (human CD34⁺ cells, human monocytes or murine bone marrow progenitors) or culture conditions, cultured DCs did not express L-selectin after 2-3 days of culture as determined by FACS analysis with anti-L-selectin antibody (anti-mouse L-selectin from PharMingen; anti-human L-selectin from Coulter Immunotech), and could neither tether nor roll on PNAd-coated surfaces (i.e. bind selectin ligands) in the parallel plate flow chamber assay.

Example 3: Expression of E/L-Selectin Chimera Restores Selectin Ligand Binding by DCs

To restore L-selectin expression, a retroviral construct containing human L-selectin (GenBank accession number X16150), was prepared according to standard procedures. A modified murine leukemia virus was generated by transfection of the 293GPG retroviral packaging cell line (Ory et al., *Proc. Natl. Acad. Sci. USA* 93:11400-11406, 1996). The retrovirus efficiently transduced murine DCs.

Murine bone marrow derived DCs were transduced with the L-selectin-expressing retrovirus on days 2 and 4 of culture and tested at day 7 or 8. Analysis of surface L-selectin expression by antibody recognition in transduced DCs showed that very few DCs expressed L-selectin protein.

As it is known that L-selectin can be rapidly degraded from the cell surface by metalloproteinases, another retroviral construct that expressed an E/L-selectin chimera (von Andrian et al., *Cell* 82:989-999, 1995; Stein et al. *J. Exp. Med.* 189:37-49, 1999) was prepared as above. The chimeric protein contains the transmembrane and intracellular domains of L-selectin and the extracellular domain of E-selectin. DCs were transduced as described above. In contrast, 15-30% of DCs transduced with the E/L-selectin chimera expressed the chimeric protein as recognized by a monoclonal anti-human E-selectin antibody (Southern Biotechnology).

The E/L-selectin transduced DCs were able to tether and roll both *in vitro* and *in vivo* on PNAd, as observed with the parallel plate flow chamber assay and using intravital microscopy, respectively.

Example 4: DC Binding to PNAd is Restored by Activated Platelets

Next it was determined whether DCs can bind activated platelets and if so whether the DC-platelet could tether and roll on PNAd. Platelets were activated with thrombin receptor activating protein (TRAP; Bachem, PA; 25 μ m for 5 min) or thrombin (Sigma Chemical Co., St. Louis, MO; 0.5 U/ml for 5 min) according to standard procedures (see, e.g., Diacovo et al., 1996). Fluorescently-labeled activated platelets (murine and human) were incubated with fluorescently-labeled murine DCs and binding was observed with a fluorescence activated cell sorter and confocal microscopy. DCs readily bound to activated platelets. Preincubation of cultured DCs with activated platelets also conferred the capacity to tether and roll on PNAd in the parallel plate flow chamber assay. Microscopy confirmed that activated platelets were bound to the rolling DCs. Activated platelets also mediate rolling of DCs in mouse high endothelial venules *in vivo* as observed by intravital microscopy.

Example 5: Localization of selectin expression on dendritic cells

Murine bone marrow-derived dendritic cells were transformed with the E/L-selectin-expressing CMMP retrovirus as described above and then stained with mouse-anti-human E-

selectin monoclonal antibody CL37 (20 μ g/ml; mouse IgG; Dr. C. Wayne Smith, Baylor College). Subsequently, the cells were washed and stained with 12 nm colloidal gold conjugated goat-anti-mouse antibody (dilution 1:6; Jackson Labs). The cells were fixed and prepared for low-voltage scanning electron microscopy (LVSEM) and analyzed in the electron microscopy core facility at the University of Minnesota as described in by von Andrian et al. (*Cell* 82:989-999, 1995).

Fig. 1 is a digitized low-voltage scanning electron micrograph showing the surface of a murine bone marrow-derived dendritic cell at 80,000 x magnification. The length of the dotted bar within the black band at the bottom of the image is 380 nm. The clusters of bright white dots on the ridges and tips of microvilli are gold particles delineating E/L-selectin expression. A quantitative analysis of gold particle distribution on three representative cells indicates that 80 - 95% of particles are associated with microvilli, and of these, ~75% are found on the upper half and the tip. There was no detectable staining on untransformed cells and no binding was detected on transformed cells when an isotype-matched non-binding control antibody was used. The data are remarkably consistent with previous studies using E/L-selectin transfected pre-B cells (von Andrian et al., *Cell* 82:989-999, 1995).

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

All references disclosed herein are incorporated by reference.

What is claimed is:

Claims

1. A method for delivery of dendritic cells to a secondary lymphoid tissue of a subject, comprising
providing isolated genetically modified dendritic cells which express on the cell
5 surface a selectin polypeptide comprising an endothelial selectin ligand binding portion of a selectin selected from the group consisting of L-selectin, E-selectin and P-selectin, and administering the isolated genetically modified dendritic cells to the subject.
2. The method of claim 1, wherein the secondary lymphoid tissue is peripheral lymph
10 nodes.
3. The method of claim 1, wherein the secondary lymphoid tissue is appendix.
4. The method of claim 1, wherein the secondary lymphoid tissue is tonsil.
5. A method for delivery of dendritic cells to a non-lymphoid tissue of a subject where
15 selectin ligands are expressed on endothelial cells, comprising
providing isolated genetically modified dendritic cells which express on the cell
surface a selectin polypeptide comprising an endothelial selectin ligand binding portion of a
20 selectin selected from the group consisting of L-selectin, E-selectin and P-selectin, and administering the isolated genetically modified dendritic cells to the subject.
6. The method of claim 1 or 5, wherein the selectin polypeptide consists of a selectin
25 selected from the group consisting of L-selectin, E-selectin and P-selectin.
7. The method of claim 1 or 5, wherein the step of providing isolated dendritic cells
comprises isolating dendritic cells from the subject and transfecting the isolated dendritic cells
with a nucleic acid molecule which encodes the selectin polypeptide.
- 30 8. The method of claim 7, wherein the nucleic acid molecule is an expression vector.
9. The method of claim 8, wherein the expression vector is selected from the group

consisting of retroviruses, lentiviruses, adenoviruses and lambda bacteriophages.

10. The method of claim 9, wherein the expression vector is a retrovirus.

11. The method of claim 7, wherein the step of providing isolated dendritic cells further comprises culturing the isolated transfected dendritic cells to expand the isolated transfected dendritic cells.

12. The method of claim 1 or 5, wherein the step of providing isolated dendritic cells further comprises treating the isolated transfected dendritic cells with isolated activated platelets or membrane microparticles thereof which contain P selectin.

13. The method of claim 1 or 5, wherein the isolated dendritic cells are administered intravenously.

14. A method for delivery of dendritic cells to a secondary lymphoid tissue of a subject, comprising
providing isolated dendritic cells,
treating the isolated dendritic cells with isolated activated platelets or membrane microparticles thereof which contain P selectin to form platelet modified dendritic cells, and
administering the isolated platelet modified dendritic cells to the subject.

15. The method of claim 14, wherein the secondary lymphoid tissue is peripheral lymph nodes.

16. The method of claim 14, wherein the secondary lymphoid tissue is appendix.

17. The method of claim 14, wherein the secondary lymphoid tissue is tonsil.

18. A method for delivery of dendritic cells to a non-lymphoid tissue of a subject where selectin ligands are expressed on endothelial cells, comprising
providing isolated dendritic cells,

treating the isolated dendritic cells with isolated activated platelets or membrane microparticles thereof which contain P selectin to form platelet modified dendritic cells, and administering the isolated platelet modified dendritic cells to the subject.

19. The method of claim 14 or 18, wherein the step of providing isolated dendritic cells further comprises culturing the isolated dendritic cells to expand the isolated dendritic cells.

20. The method of claim 14 or 18, wherein the isolated platelet modified dendritic cells are administered intravenously.

21. A method for delivery of dendritic cells to a secondary lymphoid tissue of a subject, comprising

providing isolated dendritic cells,

providing isolated activated platelets or membrane microparticles thereof which

contain P selectin,

administering the isolated dendritic cells and the isolated activated platelets or membrane microparticles thereof to the subject, wherein the isolated activated platelets or membrane microparticles thereof are administered prior to or concurrently with the isolated dendritic cells.

22. The method of claim 21, wherein the secondary lymphoid tissue is peripheral lymph nodes.

23. The method of claim 21, wherein the secondary lymphoid tissue is appendix.

24. The method of claim 21, wherein the secondary lymphoid tissue is tonsil.

25. A method for delivery of dendritic cells to a non-lymphoid tissue of a subject where selectin ligands are expressed on endothelial cells, comprising

providing isolated dendritic cells,

providing isolated activated platelets or membrane microparticles thereof which

contain P selectin,

administering the isolated dendritic cells and the isolated activated platelets or membrane microparticles thereof to the subject, wherein the isolated activated platelets or membrane microparticles thereof are administered prior to or concurrently with the isolated dendritic cells.

26. The method of claim 21 or 25, wherein the step of providing isolated dendritic cells further comprises culturing the isolated dendritic cells to expand the isolated dendritic cells.

27. The method of claim 21 or 25, wherein the isolated dendritic cells and the isolate activated platelets are administered intravenously.

28. A composition comprising isolated genetically modified dendritic cells which express on the cell surface a selectin polypeptide comprising an endothelial selectin ligand binding portion of a selectin selected from the group consisting of L-selectin, E-selectin and P-selectin.

29. The composition of claim 28, wherein the selectin polypeptide consists of a selectin selected from the group consisting of L-selectin, E-selectin and P-selectin.

30. The composition of claim 28, wherein the amount of the selectin polypeptide expressed on the cell surface is greater than the naturally occurring amount of the selectin [expressed on the cell surface *in vitro*] and is sufficient to target the genetically modified dendritic cells to peripheral lymph nodes.

31. The composition of claim 28, wherein the isolated dendritic cells are transfected with a nucleic acid molecule which encodes the selectin polypeptide.

32. The composition of claim 31, wherein the nucleic acid molecule is an expression vector.

33. The composition of claim 32, wherein the expression vector is selected from the group consisting of retroviruses, lentiviruses, adenoviruses and lambda bacteriophages.

34. The composition of claim 33, wherein the expression vector is a retrovirus.

35. The composition of claim 28, further comprising isolated activated platelets or membrane microparticles thereof which contain P selectin.

36. A composition comprising isolated dendritic cells and isolated activated platelets or membrane microparticles thereof which contain P selectin.

37. A vaccine composition comprising the composition of any of claims 28-36 and an antigen.

38. The vaccine composition of claim 37, wherein the antigen is a peptide.

39. The vaccine composition of claim 37, wherein the isolated dendritic cells are loaded with the antigen *ex vivo*.

40. The vaccine composition of claim 37, wherein the isolated dendritic cells are transfected with a nucleic acid molecule which encodes the antigen.

41. The vaccine composition of claim 40, wherein the nucleic acid molecule encodes at least two antigens.

42. The vaccine composition of claim 37, wherein the antigen is selected from the group consisting of MAGE, MART, LAGE, NY-ESO-1, tyrosinase, PRAME, prostate specific antigen (PSA), BCR-ABL, T cell receptor from T cell tumors, immunoglobulins from B cell tumors, RAS and carcinoembryonic antigen (CEA).

43. The vaccine composition of claim 37, further comprising an adjuvant.

44. The vaccine composition of claim 43, wherein the adjuvant is a nucleic acid molecule encoding a polypeptide adjuvant transfected into the isolated dendritic cells

45. The vaccine composition of claim 43 or 44, wherein the adjuvant is selected from the group consisting of IL-12, GM-CSF, CD40, CD80 and CD86.

46. The vaccine composition of claim 43 or 44, wherein the adjuvant orients an immune response to lymphocyte subsets, wherein the adjuvant is selected from the group consisting of IL-4 and interferon- γ .

47. The vaccine composition of claim 43 or 44, wherein the adjuvant is selected from the group consisting of IL-10 and TGF- β .

48. A method for stimulating an immune response to an antigen in a subject comprising administering to the subject the vaccine composition of any of claims 37-46.

49. A method for inhibiting an immune response to an antigen in a subject comprising administering to the subject the vaccine composition of claim 47.

50. A method for testing the immunogenicity of an antigen, comprising providing a subject, administering to the subject the vaccine composition of any of claims 37-46, wherein the immunogenicity of the antigen is not known, and determining the immune response of the subject to the antigen.


Abstract of the Disclosure

The invention provides isolated dendritic cells genetically modified to express a selectin polypeptide, optionally treated with activated platelets or membrane microparticles thereof. The invention also provides isolated platelet modified dendritic cells. Methods for delivering the modified dendritic cells to peripheral lymph nodes and methods for using the modified dendritic cells to stimulate immune responses also are provided. Vaccine compositions containing the modified dendritic cells also are provided.

5

5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208

Fig. 1



0054063-033100